

	CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 2017 DATED 01/01/17	
JAMES SMITH CALF GROWER CATTLE FEED MEDICATED TYPE B FEED CONTAINING AMPROLIUM FORMULA #: SMITHCALFJAN17		

STATEMENT OF PURPOSE

As an aid in the prevention of coccidiosis caused by Eimeria bovis and Eimeria zuernii during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT

Amprolium	4545.45 grams/ton
	2272.73 milligrams/pound
	0.50 %

USE DIRECTIONS

Use 10 pounds of 2.5% Type B per ton of feed to produce a Type C medicated feed containing 0.0125% amprolium. Use 1,000 pounds of 2.5% Type B per ton to produce a Type C medicated feed containing 1.25% amprolium.
Feed the resulting Type C medicated feed in amounts according to the chart below to provide 227 mg amprolium per 100 lbs. of bodyweight per day for 21 days during periods of exposure of when experience indicates coccidiosis is likely to be a hazard.

Calf Weight (pounds)	amprolium required(mg)	lbs. to feed per day	Calf Weight (pounds)	amprolium required(mg)	lbs. to feed per day
100	227	0.10	300	681	0.30
150	340.5	0.15	350	794.5	0.35
200	454	0.20	400	908	0.40
250	567.5	0.25	450	1021.5	0.45

PRECAUTIONARY STATEMENTS

WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not use Amprolium in feeds containing bentonite.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE, 55555

Net Weight on Bag and/or Invoice

Label Revised 12/1/2016

	CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 2017 DATED 01/01/17	
JAMES SMITH CALF GROWER CATTLE FEED MEDICATED TYPE B FEED CONTAINING AMPROLIUM FORMULA #: SMITHCALFJAN17		

STATEMENT OF PURPOSE

As an aid in the prevention of coccidiosis caused by Eimeria bovis and Eimeria zuernii during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT

Amprolium	4545.45 grams/ton
	2272.73 milligrams/pound
	0.50 %

USE DIRECTIONS

Use 10 pounds of 2.5% Type B per ton of feed to produce a Type C medicated feed containing 0.0125% amprolium. Use 1,000 pounds of 2.5% Type B per ton to produce a Type C medicated feed containing 1.25% amprolium.
Feed the resulting Type C medicated feed in amounts according to the chart below to provide 227 mg amprolium per 100 lbs. of bodyweight per day for 21 days during periods of exposure of when experience indicates coccidiosis is likely to be a hazard.

Calf Weight (pounds)	amprolium required(mg)	lbs. to feed per day	Calf Weight (pounds)	amprolium required(mg)	lbs. to feed per day
100	227	0.10	300	681	0.30
150	340.5	0.15	350	794.5	0.35
200	454	0.20	400	908	0.40
250	567.5	0.25	450	1021.5	0.45

PRECAUTIONARY STATEMENTS

WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not use Amprolium in feeds containing bentonite.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE, 55555

Net Weight on Bag and/or Invoice

Label Revised 12/1/2016