

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 90101 DATED 01/01/17



**Star Dairy Calf CATTLE FEED
MEDICATED**

COMPLETE FEED CONTAINING DECCOX™(DECOQUINATE)

FORMULA # SD Calf Nov 16

STATEMENT OF PURPOSE

For the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuerniii* in cattle (including ruminating and non-ruminating calves and veal calves).

ACTIVE DRUG INGREDIENT

Decoquinat e 34.09 grams/ton
17.05 milligrams/pound

USE DIRECTIONS

Feed at a rate of 22.7 mg decoquinat e per 100 lb of body weight per day as indicated in the feeding chart below. Feed at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard.

Cattle WT. (pounds)	decoquinat e required (mg)	lbs. To feed per day	Cattle WT. (pounds)	decoquinat e required (mg)	lbs. to feed per day
100	22.7	1.33	600	136.2	7.99
200	45.4	2.66	700	158.9	9.32
300	68.1	4.00	800	181.6	10.65
400	90.8	5.33	900	204.3	11.99
500	113.5	6.66	1000	227	13.32

PRECAUTIONARY STATEMENTS

WARNING: Do not feed to cows producing milk for human consumption.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.

MANUFACTURED BY:

BLUE BIRD FED MILL, ANY CITY, M ANY STATE 55555

Net Weight on Bag and/or Invoice

DECCOX™ is a trademark of Alpharma Inc.

label revision 12/01/16

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 90101 DATED 01/01/17



**Star Dairy Calf CATTLE FEED
MEDICATED**

COMPLETE FEED CONTAINING DECCOX™(DECOQUINATE)

FORMULA # SD Calf Nov 16

STATEMENT OF PURPOSE

For the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuerniii* in cattle (including ruminating and non-ruminating calves and veal calves).

ACTIVE DRUG INGREDIENT

Decoquinat e 34.09 grams/ton
17.05 milligrams/pound

USE DIRECTIONS

Feed at a rate of 22.7 mg decoquinat e per 100 lb of body weight per day as indicated in the feeding chart below. Feed at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard.

Cattle WT. (pounds)	decoquinat e required (mg)	lbs. To feed per day	Cattle WT. (pounds)	decoquinat e required (mg)	lbs. To feed per day
100	22.7	1.33	600	136.2	7.99
200	45.4	2.66	700	158.9	9.32
300	68.1	4.00	800	181.6	10.65
400	90.8	5.33	900	204.3	11.99
500	113.5	6.66	1000	227	13.32

PRECAUTIONARY STATEMENTS

WARNING: Do not feed to cows producing milk for human consumption.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.

MANUFACTURED BY:

BLUE BIRD FED MILL, ANY CITY, M ANY STATE 55555

Net Weight on Bag and/or Invoice

DECCOX™ is a trademark of Alpharma Inc.

label revision 12/01/16