

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 90101 DATED 01/01/17



**Star Dairy Heifer Calf CATTLE FEED
MEDICATED**
SUPPLEMENT FEED CONTAINING DECCOX™ (DECOQUINATE)

FORMULA # SDHC Nov 16

STATEMENT OF PURPOSE

For the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in cattle (including ruminating and non-ruminating calves and veal calves).

ACTIVE DRUG INGREDIENT

Decoquinatone 113.64 grams/ton
56.82 milligrams/pound

USE DIRECTIONS

Administer as a top dress supplement or mix into the daily ration in accordance with the feeding chart below to provide 22.7 mg per 100 pounds of body weight per day. **0.40** pounds of this supplement will medicate 100 lbs. of body weight. Feed at least 28 days during periods of coccidiosis or when it is likely to be a hazard.

Cattle WT. (pounds)	decoquinatone required (mg)	lbs. to feed per day	Cattle WT. (pounds)	decoquinatone required (mg)	lbs. to feed per day
100	22.7	0.40	600	136.2	2.40
200	45.4	0.80	700	158.9	2.80
300	68.1	1.20	800	181.6	3.20
400	90.8	1.60	900	204.3	3.60
500	113.5	2.00	1000	227	4.00

PRECAUTIONARY STATEMENTS

WARNING: Do not feed to cows producing milk for human consumption.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.

MANUFACTURED BY:
BLUE BIRD FEED MILL, ANY CITY, ANY STATE

Net Weight on Bag and/or Invoice

DECCOX™ is a trademark of Alpharma Inc.

revision 12/01/16

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 90101 DATED 01/01/17



**Star Dairy Heifer Calf CATTLE FEED
MEDICATED**
SUPPLEMENT FEED CONTAINING DECCOX™ (DECOQUINATE)

FORMULA # SDHC Nov 16

STATEMENT OF PURPOSE

For the prevention of coccidiosis in cattle (including ruminating and nonruminating calves including veal calves) caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Decoquinatone 113.64 grams/ton
56.82 milligrams/pound

USE DIRECTIONS

Administer as a top dress supplement or mix into the daily ration in accordance with the feeding chart below to provide 22.7 mg per 100 pounds of body weight per day. **0.40** pounds of this supplement will medicate 100 lbs. of body weight. Feed at least 28 days during periods of coccidiosis or when it is likely to be a hazard.

Cattle WT. (pounds)	decoquinatone required (mg)	lbs. to feed per day	Cattle WT. (pounds)	decoquinatone required (mg)	lbs. to feed per day
100	22.7	0.40	600	136.2	2.40
200	45.4	0.80	700	158.9	2.80
300	68.1	1.20	800	181.6	3.20
400	90.8	1.60	900	204.3	3.60
500	113.5	2.00	1000	227	4.00

PRECAUTIONARY STATEMENTS

WARNING: Do not feed to cows producing milk for human consumption.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.

MANUFACTURED BY:
BLUE BIRD FEED MILL, ANY CITY, ANY STATE

Net Weight on Bag and/or Invoice

DECCOX™ is a trademark of Alpharma Inc.

revision 12/01/16