



CUSTOMER-FORMULA MEDICATED LABEL
 FOR INVOICE 50001 DATED 1/1/2017
Steer Feed CATTLE FEED
MEDICATED
 CONTAINS RUMENSIN™ (MONENSIN)
DO NOT FEED UNDILUTED

FORMULA # JS Steer 12-2016

STATEMENT OF PURPOSE

For increased rate of weight gain; For prevention and control of coccidiosis caused by Eimeria bovis and E. zuernii in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium) 480.00 grams/ton
 240.00 mg/pound

USE DIRECTIONS

Mixing Directions

Thoroughly mix monensin Type B medicated feed into the appropriate amount of non-medicated feed to obtain the correct concentration in the Type C medicated feed (15 to 400 g/Ton). During the first 5 days of feeding, cattle should not receive more than 100 mg/day in not less than 1 pound of feed.

Desired Level monensin g/ton (mg/lb.)*	Amount of Type B feed to add (pounds)*	Amount of nonmedicated feed (lbs.)	Feeding level of finished feed lbs./head/day
25 (12.5)	104.17	1895.83	4 to 16
100 (50)	416.67	1583.33	1 to 4
200 (100)	833.33	1166.67	1 to 2
300 (150)	1250.00	750.00	1 to 1.3
400 (200)	1666.67	333.33	1

Feeding Directions for increased rate of weight gain:

Feed to provide 50 to 200 mg of monensin per head per day in not less than 1 pound of feed. During the first 5 days of feeding cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed.

For example, a Type C feed containing 25 g/T should be fed at a rate of 4 to 16 pounds per head per day.

*Desired level of monensin in ration (25 to 400 g/ton) ÷ monensin g/ton in the Type B feed X 2000 equals the amount of the Type B feed to add per ton of nonmedicated grain and or roughage feed RUMENSIN™ is a trademark of Eli Lilly and Company

For prevention and control of coccidiosis:

Feed continuously to provide 0.14 to 0.42 milligrams of monensin per pound of bodyweight per day, depending upon the severity of challenge, up to 200 mg monensin per head per day. Refer to the chart below for examples.

ANIMAL WEIGHT (POUNDS)	MONENSIN MG PER DAY MINIMUM	MONENSIN MG PER DAY MAXIMUM
200	28.0	84.0
300	42.0	126.0
400	56.0	168.0
500	70.0	200.0
600	84.0	200.0
700	98.0	200.0
800	112.0	200.0

Desired Intake of Monensin mg/hd/day	Type C Feed Monensin level g/Ton (mg/lb.)	Recommended Feeding Rate (lbs./hd/day)
60	15 (7.5)	8
60	30 (15)	4
60	60 (30)	2
100	100 (50)	2
200	200 (100)	2
150	300 (150)	1
200	400 (200)	1

Caution: Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555
 Net Weight on Bag and/or Invoice 12/20/2016



CUSTOMER-FORMULA MEDICATED LABEL
 FOR INVOICE 50001 DATED 1/1/2017
Steer Feed CATTLE FEED
MEDICATED
 CONTAINS RUMENSIN™ (MONENSIN)
DO NOT FEED UNDILUTED

FORMULA # JS Steer 12-2016

STATEMENT OF PURPOSE

For increased rate of weight gain; For prevention and control of coccidiosis caused by Eimeria bovis and E. zuernii in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium) 480.00 grams/ton
 240.00 mg/pound

USE DIRECTIONS

Mixing Directions

Thoroughly mix monensin Type B medicated feed into the appropriate amount of non-medicated feed to obtain the correct concentration in the Type C medicated feed (15 to 400 g/Ton). During the first 5 days of feeding, cattle should not receive more than 100 mg/day in not less than 1 pound of feed.

Desired Level monensin g/ton (mg/lb.)*	Amount of Type B feed to add (pounds)*	Amount of nonmedicated feed (lbs.)	Feeding level of finished feed lbs./head/day
25 (12.5)	104.17	1895.83	4 to 16
100 (50)	416.67	1583.33	1 to 4
200 (100)	833.33	1166.67	1 to 2
300 (150)	1250.00	750.00	1 to 1.3
400 (200)	1666.67	333.33	1

Feeding Directions for increased rate of weight gain:

Feed to provide 50 to 200 mg of monensin per head per day in not less than 1 pound of feed. During the first 5 days of feeding cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed.

For example, a Type C feed containing 25 g/T should be fed at a rate of 4 to 16 pounds per head per day.

*Desired level of monensin in ration (25 to 400 g/ton) ÷ monensin g/ton in the Type B feed X 2000 equals the amount of the Type B feed to add per ton of nonmedicated grain and or roughage feed RUMENSIN™ is a trademark of Eli Lilly and Company

For prevention and control of coccidiosis:

Feed continuously to provide 0.14 to 0.42 milligrams of monensin per pound of bodyweight per day, depending upon the severity of challenge, up to 200 mg monensin per head per day. Refer to the chart below for examples.

ANIMAL WEIGHT (POUNDS)	MONENSIN MG PER DAY MINIMUM	MONENSIN MG PER DAY MAXIMUM
200	28.0	84.0
300	42.0	126.0
400	56.0	168.0
500	70.0	200.0
600	84.0	200.0
700	98.0	200.0
800	112.0	200.0

Desired Intake of Monensin mg/hd/day	Type C Feed Monensin level g/Ton (mg/lb.)	Recommended Feeding Rate (lbs./hd/day)
60	15 (7.5)	8
60	30 (15)	4
60	60 (30)	2
100	100 (50)	2
200	200 (100)	2
150	300 (150)	1
200	400 (200)	1

Caution: Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555
 Net Weight on Bag and/or Invoice 12/20/2016