

|  |   |   |             |                |  |                       |
|--|---|---|-------------|----------------|--|-----------------------|
|   | <b>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT</b><br>FOR INVOICE 789456 DATED 01/01/17 |  |             |                |  |                       |
| <b>SMITH Finish Racto7.2 SWINE FEED<br/>MEDICATED</b>  |   |   |             |                |  |                       |
| TYPE C SWINE FEED CONTAINING RACTOPAMINE<br><br>FORMULA #: SMITHSWINEJan17   |   |   |             |                |  |                       |
| <b>STATEMENT OF PURPOSE</b>  |   |   |             |                |  |                       |
| For increased rate of weight gain, improved feed efficiency, and increased carcass leanness in finishing swine, weighing not less than 150 pounds, fed a complete ration containing at least 16% crude protein for the last 45 - 90 pounds of gain prior to slaughter. |   |   |             |                |  |                       |
| <b>ACTIVE DRUG INGREDIENT</b>  |   |   |             |                |  |                       |
| <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 70%;">Ractopamine</td> <td style="text-align: right;">7.20 grams/ton</td> </tr> <tr> <td></td> <td style="text-align: right;">3.60 milligrams/pound</td> </tr> </table>                 |   |   | Ractopamine | 7.20 grams/ton |  | 3.60 milligrams/pound |
| Ractopamine  | 7.20 grams/ton  |   |             |                |  |                       |
|  | 3.60 milligrams/pound   |   |             |                |  |                       |
| <b>USE DIRECTIONS</b>  |   |   |             |                |  |                       |
| For finishing swine, weighing not less than 150 pounds, fed a complete ration containing at least 16% crude protein for the last 45 to 90 lbs of gain prior to slaughter. Feed continuously as sole ration.  |   |   |             |                |  |                       |
| <b>PRECAUTIONARY STATEMENTS</b>  |   |   |             |                |  |                       |
| <b>CAUTION:</b> Ractopamine may increase the number of injured and/or fatigued pigs during marketing. Not for use in breeder swine. No increased benefit has been shown when ractopamine concentrations in the diet are greater than 4.5 g/ton.                        |   |   |             |                |  |                       |
| MANUFACTURED BY:<br>BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555<br><br>Net Weight on Bag and/or Invoice   |   |   |             |                |  |                       |
| revision 12/01/16  |   |   |             |                |  |                       |

|  |   |   |             |                |  |                       |
|--|---|---|-------------|----------------|--|-----------------------|
|   | <b>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT</b><br>FOR INVOICE 789456 DATED 01/01/17 |  |             |                |  |                       |
| <b>SMITH Finish Racto7.2 SWINE FEED<br/>MEDICATED</b>  |   |   |             |                |  |                       |
| TYPE C SWINE FEED CONTAINING RACTOPAMINE<br><br>FORMULA #: SMITHSWINEJan17   |   |   |             |                |  |                       |
| <b>STATEMENT OF PURPOSE</b>  |   |   |             |                |  |                       |
| For increased rate of weight gain, improved feed efficiency, and increased carcass leanness in finishing swine, weighing not less than 150 pounds, fed a complete ration containing at least 16% crude protein for the last 45 - 90 pounds of gain prior to slaughter. |   |   |             |                |  |                       |
| <b>ACTIVE DRUG INGREDIENT</b>  |   |   |             |                |  |                       |
| <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 70%;">Ractopamine</td> <td style="text-align: right;">7.20 grams/ton</td> </tr> <tr> <td></td> <td style="text-align: right;">3.60 milligrams/pound</td> </tr> </table>                 |   |   | Ractopamine | 7.20 grams/ton |  | 3.60 milligrams/pound |
| Ractopamine  | 7.20 grams/ton  |   |             |                |  |                       |
|  | 3.60 milligrams/pound   |   |             |                |  |                       |
| <b>USE DIRECTIONS</b>  |   |   |             |                |  |                       |
| For finishing swine, weighing not less than 150 pounds, fed a complete ration containing at least 16% crude protein for the last 45 to 90 lbs of gain prior to slaughter. Feed continuously as sole ration.  |   |   |             |                |  |                       |
| <b>PRECAUTIONARY STATEMENTS</b>  |   |   |             |                |  |                       |
| <b>CAUTION:</b> Ractopamine may increase the number of injured and/or fatigued pigs during marketing. Not for use in breeder swine. No increased benefit has been shown when ractopamine concentrations in the diet are greater than 4.5 g/ton.                        |   |   |             |                |  |                       |
| MANUFACTURED BY:<br>BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555<br><br>Net Weight on Bag and/or Invoice   |   |   |             |                |  |                       |
| revision 12/01/16  |   |   |             |                |  |                       |