TAKE TIME
\odot
OBSERVE LAB

CUSTOMER -FORMULA LABEL ATTACHMENT FOR INVOICE 2017 DATED 1/1/2017

Calf Feed (Summer) CATTLE FEED CONTAINS ALTOSID[™]



FORMULA # JSMITHCOWJAN17

STATEMENT OF PURPOSE

To prevent the breeding of horn flies in the manure of treated cattle.

This product prevents the emergence of adult horn flies from the manure of the treated cattle. Begin use in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity. Introduction of the product when significant numbers of adult flies are present requires the use of adulticide sprays or dusts.

ACTIVE PEST	ICIDE INGRE	EDIENT	
S-Methoprene (Altosid®)		90.91 grams/ton	
	equals	45.45 mg/lb	
	equals	0.0100 %	
USE D	DIRECTIONS		
For continuous feed	ding during th	ne fly season	

When fed as free-choice fed mineral:

Place the mixture of Altosid Premix and minerals near watering or loafing areas and provide 1 feeder for each 15 to 20 head. Put out only a 5 to 7 day supply of the mixture at one time and protect it from rain.

When the recommended mixture of Altosid premix and mineral is consumed at the rate of 0.2 oz. (0.0125 lb) or 0.4 (0.025 lb) oz. per 100 lbs. of body weight per day, which is equivalent to 2 oz. or 4 oz, for a 1000 lb, animal per day, this rate will supply the recommended average daily intake of 1.13 mg of S-Methoprene per 100 lbs of body weight per day.

When fed in a mixed-ration feed:

Provide 1.13 mg of S-Methoprene per 100 lbs, of body weight per day as indicated in the feeding schedule below

Cattle Weight	S-Methoprene	lbs. To feed	Cattle Weight	S-Methoprene	lbs. To feed
(pounds)	required (mg)	per head/day	(pounds)	required (mg)	per head/day
200	2.26	0.05	900	10.17	0.22
300	3.39	0.07	1000	11.3	0.25
400	4.52	0.10	1100	12.43	0.27
500	5.65	0.12	1200	13.56	0.30
600	6.78	0.15	1300	14.69	0.32
700	7.91	0.17	1400	15.82	0.35
800	9.04	0.20	1500	16.95	0.37

PRECAUTIONARY STATEMENTS

CAUTION-KEEP OUT OF REACH OF CHILDREN

See the attached

Altosid® Premix 0.4% label for information related to safety (i.e., Personal protective equip., first aid environmental hazards etc.

> MANUFACTURED BY: BLUE BIRD FEED MILL. ANY CITY, ANY STATE 55555

Net Weight on Bag and/or Invoice ALTOSID[™] is a trademark of Wellmark International

revision 12/1/2016

IME	CU	STOMER -FOR	MULA LABEL ATTACHN	/ENT
D	FOR INVOICE	2017	DATED	1/1/
E LABEL		Calf Feed	(Summer) CATTLE FE	ED
UNS		CONT	AINS AI TOSID [™]	

DATED 1/1/2017 ner) CATTLE FEED CONTAINS ALTOSID[™]



FORMULA # JSMITHCOWJAN17

Œ

STATEMENT OF PURPOSE

To prevent the breeding of horn flies in the manure of treated cattle.

This product prevents the emergence of adult horn flies from the manure of the treated cattle. Begin use in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity. Introduction of the product when significant numbers of adult flies are present requires the use of adulticide sprays or dusts.

ACTIVE PESTICIDE INGREDIENT				
	90.91 grams/to	on		
equals	45.45 mg/lb			
equals	0.0100 %			
IRECTION	S			
For continuous feeding during the fly season				
	equals equals DIRECTION	90.91 grams/to equals 45.45 mg/lb equals 0.0100 % IRECTIONS		

When fed as free-choice fed mineral:

Place the mixture of Altosid Premix and minerals near watering or loafing areas and provide 1 feeder for each 15 to 20 head. Put out only a 5 to 7 day supply of the mixture at one time and protect it from rain.

When the recommended mixture of Altosid premix and mineral is consumed at the rate of 0.2 oz. (0.0125 lb) or 0.4 (0.025 lb) oz. per 100 lbs. of body weight per day, which is equivalent to 2 oz. or 4 oz, for a 1000 lb, animal per day, this rate will supply the recommended average daily intake of 1.13 mg of S-Methoprene per 100 lbs of body weight per day.

When fed in a mixed-ration feed:

Provide 1.13 mg of S-Methoprene per 100 lbs, of body weight per day as indicated in the feeding schedule below

Cattle Weight	S-Methoprene	lbs. To feed	Cattle Weight	S-Methoprene	lbs. To feed
(pounds)	required (mg)	per head/day	(pounds)	required (mg)	per head/day
200	2.26	0.05	900	10.17	0.22
300	3.39	0.07	1000	11.3	0.25
400	4.52	0.10	1100	12.43	0.27
500	5.65	0.12	1200	13.56	0.30
600	6.78	0.15	1300	14.69	0.32
700	7.91	0.17	1400	15.82	0.35
800	9.04	0.20	1500	16.95	0.37

PRECAUTIONARY STATEMENTS

CAUTION-KEEP OUT OF REACH OF CHILDREN

Altosid® Premix 0.4% label for information See the attached related to safety (i.e.. Personal protective equip., first aid environmental hazards etc.

> MANUFACTURED BY: BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Net Weight on Bag and/or Invoice ALTOSID[™] is a trademark of Wellmark International

revision 12/1/2016